



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$= \frac{1}{2}(2-e^2) \int_0^{\frac{\pi}{2}} dx \sqrt{1 - \frac{e^4}{(2-e^2)^2} \sin^2 \lambda} = \frac{1}{2}(2-e^2) E\left(\frac{e^2}{2-e^2}, \frac{\pi}{2}\right) = \frac{1}{2}, \text{ if } e=1.$$

Also solved by OTTO GEUKELER, WILLIAM HOOVER, ARTEMAS MARTIN, F. P. MATZ, and G. B. M. ZERR.

21. Proposed by T. JOHN COLE, Columbus, Ohio.

In the equilateral triangle ABC , AB the base is 10 feet. With B as a center an arc is drawn from C to A ; likewise with A as a center an arc is drawn from C to B . What is the volume of the solid generated by revolving the figure about the altitude of the triangle as an axis.

I. Solution by H. C. WHITAKER, B. S., C. E., Professor of Mathematics, Manual Training School, Philadelphia, Pennsylvania.

The triangle revolving describes a cone the volume of which is $\frac{1}{3}\pi \cdot 25 \cdot 5\sqrt{3} = 226.725$.

The area of either segment is $50\left(\frac{\pi}{3} - \frac{\sqrt{3}}{2}\right) = 9.05861$; the distance of the center of gravity of the segment from the center of the circle $= \frac{1000}{12 \text{ Area}}$, and from the axis of revolution is $\left(\frac{1000}{12 \text{ Area}} - 5 \sec 30^\circ\right) \cos 30^\circ = 2.9669$.

The product of the area by the path of the center of gravity $9.05861 \times 2 \cdot \pi \cdot 2.9669 = 168.865$.

Adding to this the volume of the cone, the answer is 395.59 cubic feet.

II. Solution by JOHN DOLMAN, Jr., Counsellor at Law, Philadelphia Pennsylvania.

Let the origin be at the centre of the base. Put the side of the triangle $= 2r$, and let y = the radius of any circular horizontal section at any variable height x .

Then $V = \int_0^{r\sqrt{3}} \pi y^2 dx \dots (1)$. The equation of the arc forming the side is $4r^2 = x^2 + (y+r)^2$ from which $y^2 = 5r^2 - x^2 - 2r\sqrt{(4r^2 - x^2)}$ which value of y^2 substituted in (1) gives $V = \pi \int_0^{r\sqrt{3}} [(5r^2 - x^2 - 2r\sqrt{(4r^2 - x^2)})] dx$

$$= \pi \left[5r^2 x - \frac{1}{3} x^3 - r [x\sqrt{(4r^2 - x^2)} + 4r^2 \sin^{-1} \frac{x}{2r}] \right]_0^{r\sqrt{3}}$$

$= \pi(3r^3\sqrt{3} - \frac{4}{3}\pi r^3) = \frac{\pi r^3}{3}(9\sqrt{3} - 4\pi) = 3.164722r^3$, and putting $r=5$ this gives $V = 395.59 +$ cubic feet.

III. Solution by OTTO GEUKELER, Bloomington, Indiana.

The equation of circle with O as origin is $y^2 + (x-5)^2 = 100$, or $x=5$


$-\sqrt{(100-y^2)}$ on part of curve in question. Then from conditions of problem

$$V=2\pi \int_0^{5\sqrt{3}} \int_0^{\sqrt{100-y^2}} x dx dy = \pi \int_0^{5\sqrt{3}} [125-y^2-10] (a^2-y^2) dy.$$

$$= \pi \left[125 \times 5\sqrt{3} - 125\sqrt{3} - 1000 \int_0^{\theta=\sin^{-1} \frac{1}{\sqrt{3}}} \cos^2 \theta d\theta \right]$$

$$= \pi \left[125 \times 4 \times \sqrt{3} - 1000 \left(\frac{1}{8} + \frac{\pi}{6} \right) \right] = \pi \left[375\sqrt{3} - \right.$$

$$\left. \frac{1000}{6} \pi \right] = \frac{125\pi}{3} (9\sqrt{3} - 4\pi) = 395.59 + \text{cubic feet.}$$



IV. Solution by W. WIGGINS, Richmond, Indiana.

The volume generated is equal to that of cylinder of base CMA , and altitude equal to the length of the path sculpt out by R , the center of gravity of the area CMA . (Guldins theorem)

Taking B as the origin of co-ordinates and BA as the axis of x , we have for the area of AMC ,

$$\text{Area} = \frac{1}{2} \cdot 100 \cdot \frac{\pi}{3} - \frac{1}{2} \cdot 100 \cdot \frac{1}{2} \cdot \frac{1}{2} \sqrt{3} = \frac{1}{6} (100\pi - 75\sqrt{3}).$$

If x', y' denote the co-ordinates of c, g , of the area CMA , x' . area $CMA = \int_0^{10} x' \sqrt{(100-x^2)} dx$.

$$x'. \text{ area} = \frac{1}{3} (100-x^2)^{\frac{3}{2}} \quad x'. \text{ area} = \frac{1}{3} y^3, \text{ where } y = 5\sqrt{3}.$$

$$\therefore x' = \frac{[5\sqrt{3}]^3}{3 \times \text{Area}}, \text{ or } x' = \frac{125\sqrt{3}}{\text{Area}}, \therefore SR = \frac{125\sqrt{3}}{\text{Area}} - 5. \text{ Therefore if}$$

h denote the altitude of the equivalent cylinder, we have $h = 2\pi \left(\frac{125\sqrt{3}}{\text{Area}} - 5 \right)$.

$$V = 2\pi \left(\frac{125\sqrt{3}}{\text{Area}} - 5 \right) \times \text{Area} = 2\pi (125\sqrt{3} - 5 \text{ Area}) = 2\pi [125\sqrt{3}$$

$$- \frac{1}{6} (100\pi - 75\sqrt{3})] = \frac{125\pi}{3} (9\sqrt{3} - 4\pi) = 395.59 + \text{cubic feet.}$$

Also variously solved by C. W. M. BLACK, A. L. FOOTE, J. F. W. SCHEFFER, G. B. M. ZERR, J. H. DRUMMOND, F. P. MATZ, and P. H. PHILBRICK.

PROBLEMS.

28. Proposed by F. P. MATZ, M. Sc., Ph. D., Professor of Mathematics and Astronomy in New Windsor College, New Windsor, Maryland.

How far from the stage must Miss Love sit in order that she may see to best advantage Mr. Rich deliver the valedictory oration?

29. Proposed by CHARLES E. MYERS, Canton, Ohio.

A hen running at the rate of $n=2$ feet per second, on the circumference of a circle, radius $sr=50$ feet, is observed by a hawk $a=600$ feet directly above the center.